espacenet - Bibliographic data Page 1 of 1



## Bibliographic data: JP 2004515201 (A)

POLYPEPTIDES HOMOLOGOUS TO VEGF AND BMP1

Publication 2004-05-27 date:

Inventor(s):

Applicant(s):

Classification: international:

A61K3800, A61K4500, A61P1702; A61P1908, A61P400; C07K1482; C07K1624; C07K1900; C12W176; C12W1519; C12W1721; C12W1509; C12W1519; C12W1519; C12W510; C12P2102; C12P2108; C12Q168; G01W3376; G01W3376; G01W33757; G01W3368; [G01-7]; A61K4500; A61P1702; A61P1908; A61P900; C07K14/52; C07K16/24; C07K1900; C12W115; C12W1199; C12W151; C12W1509;

C12N5/10; C12P21/02; C12P21/08; C12Q1/68; G01N33/15; G01N33/50; G01N33/577; G01N33/68

- European: C07K14/52

Application JP20000536860T 19990310

Priority number(s): US19980040220 19980317; US19980184216 19981102; WO1999US05190 19990310

JP 4532733 (B2)

WO 9947677 (A2)
WO 9947677 (A3)
WO 9947677 (A3)

published as: • US 2008095773 (A1) • US 2010331251 (A1)

more

Abstract not available for JP 2004515201 (A) Abstract of corresponding document: WO 9947677 (A2)

The present investion involves the identification and proparation of vascular endothelial growth factor-E\_(VEG-F\_E) VEG-F\_E is a navel polyappide related to vasoviar enriphibility growth factor (VEG-F) and bone morphopenetic protein 1, VEG-F\_E has homology to VEG-F including conservation of the amino acids required for activity of VEG-F. VEG-F\_E can be useful in wound regular, as well as in the generation and regeneration of tissue.

Last updated, 26 04 2011 Worldwide Database 5.7 23; 93p